

There is a class of problems in calculus, known as related rate problems. Here's the idea. You know the rate of change (often with respect to time) of one quantity, such as the volume of a spherical balloon. You want to know the rate of change of some other related quantity (e.g. the radius of the balloon). Here are the steps you take to solve a problem like this:


1. Identify the quantity you already know a rate of change of (say,  $V$ , so you know  $dV/dt$ ).
2. Identify the quantity you want a rate of change of (say,  $r$ , so you want  $dr/dt$ ).
3. Find an equation that relates the two quantities ( $V$  and  $r$ ). This can be the hard part. Drawing a picture can help.
4. Now take a derivative with respect to  $t$  of both sides of the equation, treating both  $V$  and  $r$  as functions of  $t$ .
5. Substitute all known data into the result (typically  $V$ ,  $r$  and  $dV/dt$ ) to determine  $dr/dt$ .

We'll repeat this procedure with a bunch of examples.

1. Air is being pumped into a spherical balloon so that its volume increases at a rate of 4.5  $\text{ft}^3/\text{min}$ . How fast is the radius of the balloon increasing when the diameter is 4 ft?

0)  $t \rightarrow$  time, in minutes

1)  $V \rightarrow$  volume of balloon, in  $\text{ft}^3$   
 know  $\frac{dV}{dt} = \frac{9}{2} \text{ft}^3/\text{minute}$



2)  $r \rightarrow$  radius of balloon, in ft  
 want  $\frac{dr}{dt}$

3)  $V = \frac{4}{3}\pi r^3$

4)  $\frac{dV}{dt} = 4\pi r^2 \frac{dr}{dt}$

5)  $\frac{dr}{dt} = \frac{1}{4\pi r^2} \frac{dV}{dt}$   
 at  $r = 2 \text{ft}$  ( $d = 4 \text{ft}$ )  
 $\frac{dr}{dt} = \frac{1}{4\pi \cdot 4} \cdot \frac{9}{2} = \frac{9}{32\pi} \text{ft}^3/\text{min}$

2. Water runs into a conical tank at the rate of  $9 \text{ ft}^3/\text{min}$ . The tank stands point down and has a height of 10 ft and a base radius of 5 ft. How fast is the water level rising when the water is 6 ft deep?

0)  $t \rightarrow$  time, in minutes

1)  $V \rightarrow$  volume of water in tank, in  $\text{ft}^3$

know  $\frac{dV}{dt} = 9 \text{ ft}^3/\text{min}$


2)  $h \rightarrow$  height of water in tank, in ft

want  $\frac{dh}{dt}$

3)  $r \rightarrow$  radius of top circle of water, in ft.

$$V = \frac{1}{3} \pi r^2 h$$

Need to relate  $r$  to  $h$ :  $\frac{r}{h} = \frac{5}{10}$  by similar  $\Delta$ 's.



$$V = \frac{\pi}{12} h^3$$

$$4) \frac{dV}{dt} = \frac{3\pi}{12} h^2 \frac{dh}{dt}$$

$$5) \frac{dh}{dt} = \frac{4}{\pi h^2} \frac{dV}{dt}$$

$$= \frac{4}{\pi 6^2} \cdot 9 = \frac{1}{\pi} \text{ ft/min}$$

3. A street light is mounted at the top of a 10-ft-tall pole. A woman 5 ft tall walks away from the pole along a straight path at a speed of 5 ft/s. How fast is the tip of her shadow moving when she is 40 ft from the pole?


0)  $t$ , time in seconds

1)  $x \rightarrow$  distance of woman from pole in ft

know:  $\frac{dx}{dt} = 5 \text{ ft/sec}$

2)  $l \rightarrow$  distance of shadow tip from pole, in ft.

want  $\frac{dl}{dt}$



3) similar  $\Delta$ 's:  $\frac{l-x}{5} = \frac{l}{10}$

$$2l - 2x = l$$

$$\text{so } l = 2x$$

$$4) \frac{dl}{dt} = 2 \frac{dx}{dt}$$

$$5) \frac{dl}{dt} = 2 \cdot 5 \text{ ft/sec}$$

$$= 10 \text{ ft/sec}$$

(the at 40 ft part is not needed!)

4. A pebble dropped into a calm pond, causing ripples in the form of circles. The radius  $r$  of the outer ripple is increasing at a constant rate of 1 foot per second. When the radius is 4 feet, at what rate is the area  $A$  of the water disturbed changing?

0)  $t$ , time in seconds

1)  $r \rightarrow$  radius of circle of disturbed water, in ft

know:  $\frac{dr}{dt} = 1 \text{ ft/sec}$

2)  $A \rightarrow$  area of circle of disturbed water, in  $\text{ft}^2$

want  $\frac{dA}{dt}$

3)  $A = \pi r^2$

4)  $\frac{dA}{dt} = 2\pi r \frac{dr}{dt}$

5) at  $r = 4 \text{ ft}$ :

$$\frac{dA}{dt} = 2\pi \cdot 4 \cdot 1 = 8\pi \text{ ft}^2/\text{sec}$$



5. A hot air balloon rising straight up from a level field is tracked by a range finder 500 feet from the lift-off point. At the moment the range finder's elevation angle is  $\pi/4$ , the angle is increasing at the rate of 0.14 radians/min. How fast is the balloon rising at that moment?

0)  $t \rightarrow$  time in minutes

1)  $\theta$ : elevation angle, radians


know  $\frac{d\theta}{dt} = 0.14 \text{ rad/min}$

2)  $H$ : height of balloon in ft

want  $\frac{dH}{dt}$

3)  $\tan \theta = \frac{H}{500}$

4)  $\sec^2 \theta \frac{d\theta}{dt} = \frac{1}{500} \frac{dH}{dt}$



5)  $\frac{dH}{dt} = 500 \sec^2 \theta \frac{d\theta}{dt}$

At  $\theta = \pi/4$ ,  $\sec \theta = \frac{1}{\cos \theta} = \sqrt{2}$

$$\frac{dH}{dt} = 1000 (0.14) = 140 \text{ ft/min}$$

6. The standard 12 foot ladder rests against a vertical wall. If the bottom of the ladder slides away from the wall at a rate of 1ft/s, how fast is the top of the ladder sliding down the wall when the bottom of the ladder is 6 ft from the wall?

0)  $t$ : time in seconds

1)  $x$ : distance of base of ladder from wall, in ft

know  $\frac{dx}{dt} = 1 \text{ ft/sec}$

2)  $y$ : height of ladder against wall, in ft.

want  $\frac{dy}{dt}$


3)  $x^2 + y^2 = 12^2$

4)  $2x \frac{dx}{dt} + 2y \frac{dy}{dt} = 0$

5)  $\frac{dy}{dt} = -\frac{x}{y} \frac{dx}{dt}$

$x=6, y = \sqrt{12^2 - 6^2} = \sqrt{108}$

$\frac{dy}{dt} = -\frac{6}{\sqrt{108}} \cdot 1 = \boxed{-\frac{6}{\sqrt{108}} \text{ ft/s}}$



7. A police cruiser, approaching a right-angled intersection from the north, is chasing a speeding car that has turned the corner and is now moving straight east. When the cruiser is 0.6 mi north of the intersection and the car is 0.8 mi to the east, the police determine that the distance between them and the car they are chasing is increasing at a rate of 20 mph. If the cruiser is moving at 60 mph at the instant of measurement, what is the speed of the car? [Hint: You'll need to relate three quantities here!]

0)  $t$ , time in hours


two known rates of change!

1)  $y$ : distance of police from intersection, in miles

know  $\frac{dy}{dt} = -60 \text{ mph}$

$l$ : distance of police from other car, in miles

know  $\frac{dl}{dt} = 20 \text{ mph}$



2)  $x$ : distance of other car from intersection, in miles

want  $\frac{dx}{dt}$

3)  $x^2 + y^2 = l^2$

5)  $\frac{dx}{dt} = \frac{1}{x} \left[ l \frac{dl}{dt} - y \frac{dy}{dt} \right]$

$x = \frac{8}{10}, y = \frac{6}{10}, l^2 = \frac{8^2 + 6^2}{100} = \frac{100}{100} = 1$

4  $\Rightarrow l = 1$

$\frac{dx}{dt} = \frac{10}{8} \left[ 1 \cdot 20 - \frac{6}{10} (-60) \right] = \frac{10}{8} \cdot 56 = \boxed{70 \text{ mph}}$