


Name: _____

_____ / 25

Circle one: Rhodes (F01) | Bueller (F02)

25 points possible. No aids (book, calculator, etc.) are permitted. Show all work and use proper notation for full credit.

1. [6 points] The graph of f is shown. Evaluate each integral by interpreting it in terms of areas.



a. $\int_{-2}^0 f(x) dx =$


b. $\int_4^0 f(x) dx =$

c. $\int_{-2}^4 f(x) dx =$

2. [6 points] A particle is moving with the given acceleration $a(t)$ and other data. Find the position $s(t)$ of the particle.

$$a(t) = 3 \cos t - 2 \sin t, \quad s(0) = 0, \quad v(0) = 4$$

3. [8 points] Consider the graph of $f(x) = \frac{1}{1+x}$ below.



- a. In the figure above, sketch four rectangles corresponding to the $n = 4$ Riemann sum on the interval $1 \leq x \leq 3$. Use left endpoints.
- b. Compute the numerical value of the Riemann sum illustrated in part a. Express your answer as a single fraction.

- c. Is your numerical value in part b an overestimate or an underestimate of $\int_1^3 \frac{1}{1+x} dx$?

4. [5 points] Use the Midpoint Rule with $n = 2$ subintervals to approximate the integral. Express your answer as a single fraction.

$$\int_0^4 x 2^{-x} dx \approx$$